

Radiation and Dynamics of Dust Particle

J. Klačka

Astronomical Institute, Faculty of Mathematics, Physics, and Informatics

Comenius University, Mlynská dolina, 842 48 Bratislava, Slovak Republic

E-mail: klacka@fmph.uniba.sk

Abstract. Relativistically covariant form of equation of motion for arbitrarily shaped dust particle (neutral in charge) under the action of electromagnetic radiation is derived – emission, scattering and absorption of radiation is considered. The result is presented in the form of optical quantities used in optics of dust particles. The obtained equation of motion represents a generalization of the Poynting-Robertson (P-R) effect, which is standardly used in orbital evolution of dust particles in astrophysics.

Simultaneous action of electromagnetic radiation and gravitational fields of the central body – star – on the motion of the particle is discussed.

Key words: relativity theory, cosmic dust

1. Introduction

The first case closely connected with relativistic equation of motion for a particle under the action of electromagnetic radiation was presented by Einstein (1905), who calculated the change of energy and light pressure for plane mirror. Later on, Robertson (1937) has presented equation of motion for perfectly absorbing spherical particle – this case has been applied to astronomical situations for several decades as the well-known Poynting-Robertson (P-R) effect.

It is inevitable to have in disposal general equation of motion. Equation of motion for all types of particles, not only for the special cases treated by the above cited authors.

Real particles scatter radiation in a more complicated manner and particles of various optical properties occur. Moreover, thermal emission has to be included, in order to be the effect of radiation complete. Thus, this paper generalizes the results presented in Klačka (2000a, 2000b, 2001, 2002a) and Klačka and Kocifaj (2001), in which also results of detailed numerical calculations for (sub)micron sized dust particles in the Solar System may be found – motion of real particles may significantly differ from that corresponding to the P-R effect.

2. Proper reference frame of the particle – stationary particle

The term “stationary particle” will denote particle which does not move in a given inertial frame of reference. Primed quantities will denote quantities measured in the proper reference frame of the particle.

The flux density of photons scattered into an element of solid angle $d\Omega' = \sin \theta' d\theta' d\phi'$ is proportional to $p'(\theta', \phi') d\Omega'$, where $p'(\theta', \phi')$ is “phase function”. Phase function depends on orientation of the particle with respect to the direction of the incident radiation and on the particle characteristics; angles θ' , ϕ' correspond to the direction (and orientation) of travel of the scattered radiation, θ' is polar angle and it equals zero for the case of the travel of the ray in the orientation identical with the unit vector $\mathbf{e}'_1$ of the incident radiation. The phase function fulfills the condition

$$\int_{4\pi} p'(\theta', \phi') d\Omega' = 1 . \quad (1)$$

The momentum of the incident beam of photons which is lost in the process of its interaction with the particle is proportional to the cross section C'_{ext} (extinction). The part proportional to C'_{abs} (absorption) is completely lost and the part proportional to $C'_{ext} - C'_{abs} = C'_{sca}$ (scattering) is again reemitted. The differential scattering cross section $dC'_{sca}/d\Omega' \equiv C'_{sca} p'(\theta', \phi')$ depends on the polarization state of the incident light as well as on the incidence and scattering directions (e. g., Mishchenko et al. 2002).

The momentum (per unit time) of the scattered photons into an element of solid angle $d\Omega'$ is

$$d\mathbf{p}'_{sca} = \frac{1}{c} S' C'_{sca} p'(\theta', \phi') \mathbf{K}' d\Omega' , \quad (2)$$

where unit vector in the direction of scattering is

$$\mathbf{K}' = \cos \theta' \mathbf{e}'_1 + \sin \theta' \cos \phi' \mathbf{e}'_2 + \sin \theta' \sin \phi' \mathbf{e}'_3 . \quad (3)$$

S' is the flux density of radiation energy. The system of unit vectors used on the right-hand side of the last equation forms an orthogonal basis. The total momentum (per unit time) of the scattered photons is

$$\mathbf{p}'_{sca} = \frac{1}{c} S' C'_{sca} \int_{4\pi} p'(\theta', \phi') \mathbf{K}' d\Omega' . \quad (4)$$

The momentum (per unit time) obtained by the particle due to the interaction with radiation is

$$\frac{d\mathbf{p}'}{dt'} = \frac{1}{c} S' \left\{ C'_{ext} \mathbf{e}'_1 - C'_{sca} \int_{4\pi} p'(\theta', \phi') \mathbf{K}' d\Omega' \right\} . \quad (5)$$

As for the energy, we suppose that it is conserved: the energy (per unit time) of the incoming radiation, equals to the energy (per unit time) of the outgoing radiation (after interaction with the particle). We will use the fact that $t' = \tau$, where τ is proper time.

For the sake of brevity, we will use dimensionless efficiency factors Q'_x instead of cross sections C'_x : $C'_x = Q'_x A'$, where A' is geometrical cross section of a sphere of volume equal to the volume of the particle. Equation (5) can be rewritten to the form

$$\begin{aligned} \frac{d\mathbf{p}'}{d\tau} = \frac{1}{c} S' A' \{ [Q'_{ext} - \langle \cos \theta' \rangle Q'_{sca}] \mathbf{e}'_1 + \\ [- \langle \sin \theta' \cos \phi' \rangle Q'_{sca}] \mathbf{e}'_2 + [- \langle \sin \theta' \sin \phi' \rangle Q'_{sca}] \mathbf{e}'_3 \} . \end{aligned} \quad (6)$$

Summarizing important equations, we can write them in a short form

$$\frac{d\mathbf{p}'}{d\tau} = \frac{1}{c} S' A' \{ Q'_1 \mathbf{e}'_1 + Q'_2 \mathbf{e}'_2 + Q'_3 \mathbf{e}'_3 \} ; \quad \frac{dE'}{d\tau} = 0 . \quad (7)$$

3. Stationary frame of reference

By the term “stationary frame of reference” we mean a frame of reference in which particle moves with a velocity vector $\mathbf{v} = \mathbf{v}(t)$. The physical quantities measured in the stationary frame of reference will be denoted by unprimed symbols.

Our aim is to derive equation of motion for the particle in the stationary frame of reference. We will use the fact that we know this equation in the proper frame of reference – see Eqs. (7). We have to use Lorentz transformation for the purpose of making transformation from proper frame of reference to stationary frame of reference.

If we have a four-vector $A^\mu = (A^0, \mathbf{A})$, where A^0 is its time component and $\mathbf{A}$ is its spatial component, generalized special Lorentz transformation yields

$$\begin{aligned} A^{0'} &= \gamma (A^0 - \mathbf{v} \cdot \mathbf{A}/c) , \\ \mathbf{A}' &= \mathbf{A} + [(\gamma - 1) \mathbf{v} \cdot \mathbf{A}/v^2 - \gamma A^0/c] \mathbf{v} , \end{aligned} \quad (8)$$

where $\gamma = 1/\sqrt{1 - \mathbf{v}^2/c^2}$. The inverse generalized special Lorentz transformation is

$$\begin{aligned} A^0 &= \gamma (A^{0'} + \mathbf{v} \cdot \mathbf{A}'/c) , \\ \mathbf{A} &= \mathbf{A}' + [(\gamma - 1) \mathbf{v} \cdot \mathbf{A}'/\mathbf{v}^2 + \gamma A^{0'}/c] \mathbf{v} . \end{aligned} \quad (9)$$

Well-known four-vector is four-momentum: $p^\mu = (p^0, \mathbf{p}) \equiv (E/c, \mathbf{p})$.

3.1. Transformation of equation of motion

Eqs. (9) yield for our case represented by the second of Eqs. (7):

$$\begin{aligned} \frac{dE}{d\tau} &= \gamma \mathbf{v} \cdot \frac{d\mathbf{p}'}{d\tau} , \\ \frac{d\mathbf{p}}{d\tau} &= \frac{d\mathbf{p}'}{d\tau} + (\gamma - 1) \left(\mathbf{v} \cdot \frac{d\mathbf{p}'}{d\tau} \right) \frac{\mathbf{v}}{\mathbf{v}^2} . \end{aligned} \quad (10)$$

3.2. Flux density of radiation energy

Let us consider that a beam of incident photons is characterized by flux density of radiation energy $S' = n' h\nu' c$, or $S = n h\nu c$, where n' and n are concentrations of photons in the corresponding reference frames. The incident photon is characterized by four-momentum $p^\mu = (h\nu, h\nu \mathbf{e}_1)$. Application of Eqs. (8) to the four-momentum of the photon yields:

$$\begin{aligned} \nu' &= \nu w_1 , \\ \mathbf{e}'_1 &= \frac{1}{w_1} \{ \mathbf{e}_1 + [(\gamma - 1) \mathbf{v} \cdot \mathbf{e}_1/\mathbf{v}^2 - \gamma/c] \mathbf{v} \} , \end{aligned} \quad (11)$$

where abbreviation $w_1 \equiv \gamma (1 - \mathbf{v} \cdot \mathbf{e}_1/c)$ is used. Continuity equation $\partial_\mu j^\mu = 0$, where $j^\mu = (cn, cn\mathbf{e}_1)$ is four-vector of current density, yields $n' = nw_1$. Inserting this result together with the first of Eqs. (11) into the flux density of radiation energy S' , we obtain

$$S' = w_1^2 S . \quad (12)$$

3.3. Transformation of unit vectors

We have already found transformation of the vector $\mathbf{e}_1$. We have to found also transformations of vectors $\mathbf{e}_2$ and $\mathbf{e}_3$. Physics of the corresponding transformation is that the vectors $\mathbf{e}_2$ and $\mathbf{e}_3$ describe directions of propagation of light after interaction with the particle (see Klačka 2000a for more details). Thus, aberration of light exists for each of these unit vectors. Considerations analogous to those for vector $\mathbf{e}_1$ immediately yield

$$\begin{aligned} \mathbf{e}'_j &= \frac{1}{w_j} \{ \mathbf{e}_j + [(\gamma - 1) \mathbf{v} \cdot \mathbf{e}_j/\mathbf{v}^2 - \gamma/c] \mathbf{v} \} , \quad j = 1, 2, 3 , \\ w_j &\equiv \gamma (1 - \mathbf{v} \cdot \mathbf{e}_j/c) . \end{aligned} \quad (13)$$

3.4. Equation of motion

Inserting Eqs. (7), (12) and (13) into Eqs. (10), one easily obtains

$$\begin{aligned}\frac{dE}{d\tau} &= \sum_{j=1}^3 Q'_j w_1^2 S A' (1/w_j - \gamma) , \\ \frac{d\mathbf{p}}{d\tau} &= \sum_{j=1}^3 Q'_j \frac{w_1^2 S A'}{c^2} (c \mathbf{e}_j / w_j - \gamma \mathbf{v}) .\end{aligned}\quad (14)$$

Eqs. (14) may be rewritten in terms of four-vectors:

$$\frac{dp^\mu}{d\tau} = \frac{w_1^2 S A'}{c^2} \sum_{j=1}^3 Q'_j (c b_j^\mu - u^\mu) , \quad (15)$$

where p^μ is four-vector of the particle of mass m , $p^\mu = m u^\mu$, four-vector of the world-velocity of the particle is $u^\mu = (\gamma c, \gamma \mathbf{v})$. We have also other four-vectors

$$b_j^\mu = (1/w_j, \mathbf{e}_j / w_j) , \quad j = 1, 2, 3 . \quad (16)$$

It can be easily verified that Eq. (15) yields $dm/d\tau = 0$.

As for practical calculations we introduce:

$$\begin{aligned}b_j^0 &\equiv 1/w_j = \gamma (1 + \mathbf{v} \cdot \mathbf{e}'_j / c) , \quad j = 1, 2, 3, \\ \mathbf{b}_j &\equiv \mathbf{e}_j / w_j = \mathbf{e}'_j + [(\gamma - 1) \mathbf{v} \cdot \mathbf{e}'_j / v^2 + \gamma / c] \mathbf{v} .\end{aligned}\quad (17)$$

Within the accuracy to the first order in $\mathbf{v}/c$, Eq. (15) yields

$$\frac{d\mathbf{v}}{dt} = \frac{S A'}{m c} \sum_{j=1}^3 Q'_j [(1 - 2 \mathbf{v} \cdot \mathbf{e}_1 / c + \mathbf{v} \cdot \mathbf{e}_j / c) \mathbf{e}_j - \mathbf{v} / c] . \quad (18)$$

Eq. (15) yields as special cases the situations discussed in Einstein (1905) and Robertson (1937) – Robertson's case is obtained simply putting $Q'_1 = 1$, $Q'_2 = Q'_3 = 0$, Einstein's results require a little more calculations.

3.5. Heuristic derivation

Since we completely understand physics of Eq. (15), we are able to present short simple derivation of Eq. (18), now. We have:

- i) $d\mathbf{v}'/dt = [S' A' / (mc)] \sum_{i=1}^3 Q'_i \mathbf{e}'_i$ (see Eq. (7));
- ii) $S' = S(1 - 2\mathbf{v} \cdot \mathbf{e}_1 / c)$ (see Eq. (12)), due to the change of concentration of photons $n' = n(1 - \mathbf{v} \cdot \mathbf{e}_1 / c)$ and Doppler effect $\nu' = \nu(1 - \mathbf{v} \cdot \mathbf{e}_1 / c)$;
- iii) $\mathbf{e}'_j = (1 + \mathbf{v} \cdot \mathbf{e}_j / c) \mathbf{e}_j - \mathbf{v} / c$, $j = 1, 2, 3$ (aberration of light – see Eq. (13)).

Taking into account these physical phenomena, we finally obtain Eq. (18).

4. Thermal emission

If the particle's absolute temperature is above zero, it can emit as well as scatter and absorb electromagnetic radiation.

4.1. Proper reference frame of the particle – stationary particle

This subsection is based on the presentation by Mishchenko et. al. (2002).

The emission component of the radiation force acting on the particle of absolute temperature T' is

$$\mathbf{F}'_e(T') = -\frac{1}{c} \int_0^\infty d\omega' \int_{4\pi} \hat{\mathbf{r}}' K'_e(\hat{\mathbf{r}}', T', \omega') d\hat{\mathbf{r}}', \quad (19)$$

where the unit vector $\hat{\mathbf{r}}' = \mathbf{r}'/r'$ is given by position vector $\mathbf{r}'$ of the observation point with origin inside the particle (the emitted radiation in the far-field zone of the particle propagates in the radial direction, i. e., along the unit vector $\hat{\mathbf{r}}'$), ω' is (angular) frequency of radiation,

$$K'_e(\hat{\mathbf{r}}', T', \omega') = I'_b(T', \omega') \left\{ K'_{11}(\hat{\mathbf{r}}', \omega') - \int_{4\pi} Z'_{11}(\hat{\mathbf{r}}', \hat{\mathbf{r}}'', \omega') d\hat{\mathbf{r}}'' \right\}, \quad (20)$$

and the Planck blackbody energy distribution is given by the well-known relation

$$I'_b(T', \omega') = \frac{\hbar \omega'^3}{4 \pi^3 c^2} \left\{ \exp\left(\frac{\hbar \omega'}{k T'}\right) - 1 \right\}^{-1}. \quad (21)$$

The functions K'_{11} and Z'_{11}

$$\begin{aligned} K'_{11} &= \frac{2\pi}{k'_1} \text{Im}(S'_{11} + S'_{22}), \\ Z'_{11} &= (|S'_{11}|^2 + |S'_{12}|^2 + |S'_{21}|^2 + |S'_{22}|^2) / 2, \end{aligned} \quad (22)$$

are given by elements of the amplitude matrix $S'(\hat{\mathbf{n}}'^{sca}, \hat{\mathbf{n}}'^{inc})$:

$$\mathbf{E}'^{sca}(r' \hat{\mathbf{n}}'^{sca}) = \frac{e^{i k'_1 r'}}{r'} S'(\hat{\mathbf{n}}'^{sca}, \hat{\mathbf{n}}'^{inc}) \mathbf{E}_0'^{inc}, \quad r' \rightarrow \infty. \quad (23)$$

We assume that the incident field is a plane electromagnetic wave given by

$$\mathbf{E}'^{inc}(\mathbf{r}') = \mathbf{E}_0'^{inc} \exp\left(i k'_1 \hat{\mathbf{n}}'^{inc} \cdot \mathbf{r}'\right), \quad \omega'^2 = c^2 |\mathbf{k}'_1|^2, \quad (24)$$

since we consider vacuum as a homogeneous nonabsorbing medium. We use a couplet (ϑ', φ') , where $\vartheta' \in \langle 0, \pi \rangle$ is the polar (zenith) angle measured from the positive z' -axis and $\varphi' \in \langle 0, 2\pi \rangle$ is the azimuth angle measured from the positive x' -axis in the clockwise direction when looking in the direction of the positive z' -axis. Since the medium is assumed to be nonabsorbing, $\mathbf{E}' = E'_{\vartheta'} \hat{\boldsymbol{\vartheta}}' + E'_{\varphi'} \hat{\boldsymbol{\varphi}}'$, where $E'_{\vartheta'} \hat{\boldsymbol{\vartheta}}'$ lies in the

meridional plane (i. e., the plane through $\hat{n}'$ and the z' -axis), whereas the component $E'_{\varphi'}$, $\hat{\varphi}'$ is perpendicular to this plane; $\hat{\vartheta}'$ and $\hat{\varphi}'$ are the corresponding unit vectors such that $\hat{n}' = \hat{\vartheta}' \times \hat{\varphi}'$:

$$\begin{aligned} E_0'^{inc} &= E_{0\vartheta'}'^{inc} \hat{\vartheta}'^{inc} + E_{0\varphi'}'^{inc} \hat{\varphi}'^{inc} , \\ E'^{sca} &= E_{\vartheta'}'^{sca} \hat{\vartheta}'^{sca} + E_{\varphi'}'^{sca} \hat{\varphi}'^{sca} , \quad (\hat{r}' \cdot E'^{sca}(\mathbf{r}') = 0 \quad , \quad r' \rightarrow \infty) . \end{aligned} \quad (25)$$

4.2. Stationary frame of reference

Using orthonormal vectors $\mathbf{e}'_1$, $\mathbf{e}'_2$ and $\mathbf{e}'_3$ defined by Eq. (3), we can write Eq. (19) in the form

$$\mathbf{F}'_e(T') = \sum_{i=1}^3 [\mathbf{F}'_e(T') \cdot \mathbf{e}'_i] \mathbf{e}'_i . \quad (26)$$

Using four-vectors defined by Eq. (16), we can immediately write

$$\left(\frac{d p^\mu}{d \tau} \right)_e = \sum_{j=1}^3 Q'_{ej} b_j^\mu , \quad (27)$$

where

$$Q'_{ej} = - \mathbf{e}'_j \cdot \frac{1}{c} \int_0^\infty d\omega' \int_{4\pi} \hat{\mathbf{r}}' K'_e(\hat{\mathbf{r}}', T', \omega') d\hat{\mathbf{r}}' , \quad j = 1, 2, 3 . \quad (28)$$

As a consequence, Eqs. (16) and (27) yield

$$\left(\frac{d m}{d \tau} \right)_e = \sum_{j=1}^3 Q'_{ej} / c . \quad (29)$$

Mass of the particle decreases due to the thermal emission, alone.

5. Electromagnetic interaction – total

5.1. Proper reference frame of the particle – stationary particle

We can write, on the basis of Eqs. (7), (26) and (28):

$$\frac{d \mathbf{p}'}{d \tau} = \sum_{j=1}^3 \left(\frac{S' A'}{c} Q'_j + Q'_{ej} \right) \mathbf{e}'_j ; \quad \frac{d E'}{d \tau} = 0 . \quad (30)$$

5.2. Stationary frame of reference

On the basis of Eqs. (12), (16) and (30) we obtain

$$\frac{d p^\mu}{d \tau} = \sum_{j=1}^3 \left(\frac{w_1^2 S A'}{c} Q'_j + Q'_{ej} \right) (b_j^\mu - u^\mu / c) . \quad (31)$$

As a consequence, $d m / d \tau = 0$.

6. Gravitational and electromagnetic radiation fields

Let us consider orbital evolution of real dust particle in the Solar System, under action of gravitational and electromagnetic radiation fields of the Sun. We can write (also formulation presented in Klačka 2002b is used)

$$\begin{aligned} \frac{d\mathbf{v}}{dt} = & -\frac{GM_\odot}{r^2} \mathbf{e}_1 + \frac{GM_\odot}{r^2} \sum_{j=1}^3 \beta_j \left[\left(1 - 2 \frac{\mathbf{v} \cdot \mathbf{e}_1}{c} + \frac{\mathbf{v} \cdot \mathbf{e}_j}{c} \right) \mathbf{e}_j - \frac{\mathbf{v}}{c} \right] + \\ & \sum_{j=1}^3 Q'_{ej} \left[\left(1 + \frac{\mathbf{v} \cdot \mathbf{e}_j}{c} \right) \mathbf{e}_j - \frac{\mathbf{v}}{c} \right], \end{aligned} \quad (32)$$

where

$$\begin{aligned} \mathbf{e}_j &= (1 - \mathbf{v} \cdot \mathbf{e}'_j/c) \mathbf{e}'_j + \mathbf{v}/c, \quad j = 1, 2, 3, \\ \beta_1 &= \frac{\pi R_\odot^2}{GM_\odot m c} \int_0^\infty B_\odot(\lambda) \{C'_{ext}(\lambda/w) - C'_{sca}(\lambda/w) g'_1(\lambda/w)\} d\lambda, \\ \beta_2 &= \frac{\pi R_\odot^2}{GM_\odot m c} \int_0^\infty B_\odot(\lambda) \{-C'_{sca}(\lambda/w) g'_2(\lambda/w)\} d\lambda, \\ \beta_3 &= \frac{\pi R_\odot^2}{GM_\odot m c} \int_0^\infty B_\odot(\lambda) \{-C'_{sca}(\lambda/w) g'_3(\lambda/w)\} d\lambda, \\ w &= 1 - \mathbf{v} \cdot \mathbf{e}_1 / c, \end{aligned} \quad (33)$$

$R_\odot$ denotes the radius of the Sun and $B_\odot(\lambda)$ is the solar radiance at a wavelength of λ ; G , $M_\odot$, and r are the gravitational constant, the mass of the Sun, and the distance of the particle from the center of the Sun, respectively. The asymmetry parameter vector $\mathbf{g}'$ is defined by $\mathbf{g}' = (1/C'_{sca}) \int \mathbf{n}' (dC'_{sca}/d\Omega') d\Omega'$, where $\mathbf{n}'$ is a unit vector in the direction of scattering; $\mathbf{g}' = g'_1 \mathbf{e}'_1 + g'_2 \mathbf{e}'_2 + g'_3 \mathbf{e}'_3$, $\mathbf{e}'_1 = (1 + \mathbf{v} \cdot \mathbf{e}_1/c) \mathbf{e}_1 - \mathbf{v}/c$, $\mathbf{e}'_i \cdot \mathbf{e}'_j = \delta_{ij}$.

7. Conclusion

We have derived relativistically covariant equation of motion for dust particle under the action of electromagnetic radiation – see Eq. (31). As a generalization requires, Eq. (31) may be reduced to special cases treated by Einstein (1905), Robertson (1937) and Klačka (2000a). Application of Eq. (31) to Solar System is represented by Eqs. (32) and (33). Some other accelerations can be added to the right-hand side of Eq. (32) – e. g., gravitational perturbations of planets, solar wind effect $[(d\mathbf{v}/dt)_{s.w.} = -(\eta\beta_1 GM_\odot/(Q'_1 r^2))((\mathbf{v} \cdot \mathbf{e}_1/c)\mathbf{e}_1 + \mathbf{v}/c); \eta \approx 0.3]$, or some other nongravitational accelerations.

Acknowledgements. This paper was supported by VEGA grant No. 1/7067/20.

References

- Einstein A. 1905. Zur Elektrodynamik der bewegter Körper. *Annalen der Physik* **17**, 891-920.
- Klačka J. 2000a. Electromagnetic radiation and motion of real particle.
<http://xxx.lanl.gov/abs/astro-ph/0008510>
- Klačka J. 2000b. Aberration of light and motion of real particle.
<http://xxx.lanl.gov/abs/astro-ph/0009108>
- Klačka J. 2001. Motion of electrically neutral particle in the field of electromagnetic radiation. *Meteor Reports* **22**, 21-28.
- Klačka J. 2002a. Covariant equation of motion for a particle in an electromagnetic field.
 In: *Optics of Cosmic Dust*, G. Videen and M. Kocifaj (eds.), Kluwer Academic Publishers, Dordrecht/Boston/London, pp. 301-312.
- Klačka J. 2002b. On radiation pressure and the Poynting-Robertson effect for fluffy dust particles. <http://xxx.lanl.gov/abs/astro-ph/0208259> (submitted to *Icarus*)
- Klačka J., Kocifaj M. 2001. Motion of nonspherical dust particle under the action of electromagnetic radiation. *J. Quant. Spectrosc. Radiat. Transfer* **70/4-6**, 595-610.
- Mishchenko M., Travis L. D., Lacis A. A. 2002. Scattering, Absorption and Emission of Light by Small Particles. Cambridge University Press, Cambridge (UK), 445 pp.
- Robertson H. P. 1937. The dynamical effects of radiation in the Solar System. *Mon. Not. R. Astron. Soc.* **97**, 423-438.